




WEST NILE VIRUS

CRUDE DATA	
Number of Cases	43
Annual Incidence ^a	
LA County	0.44
California ^b	1.04
United States ^b	1.20
Age at Diagnosis	
Mean	61.5
Median	62
Range	15–94 years

^a Cases per 100,000 population.

^b Incidence calculated with 2007 population estimates from www.census.gov.



DESCRIPTION

Life Cycle and Epidemiology

West Nile virus (WNV) is a single-stranded RNA virus placed within the family Flaviviridae, genus Flavivirus. Within the genus Flavivirus, WNV has been serologically classified within the Japanese encephalitis (JE) virus antigenic complex, which includes the human pathogens JE, Murray Valley encephalitis, Saint Louis encephalitis (SLE), and Kunjin viruses.

WNV was indigenous to Africa, Asia, Europe, and Australia, and was introduced to North America in 1999, when it was first detected in New York City. The likely origin of the introduced strain was the Middle East, but the mode of introduction remains unknown. Since 1999, human and non-human WNV surveillance data has documented that WNV has extended its range through most of the continental United States as well as to Canada and Mexico.

The life cycle of the virus involves the transmission of the virus between mosquitoes and bird reservoir hosts. Humans are incidentally infected when bitten by an infected mosquito, usually a *Culex* or *Anopheles* species. The incubation period for human infection is 2 to 14 days. Birds, especially corvids such as the North American crow, are the optimal hosts for harboring and replicating the virus. Mosquitoes become infected when they feed on infected birds, which may circulate high level of viremia for several days. Infectious mosquitoes carry virus particles in their salivary glands and infect susceptible bird species during blood-meal feeding. Bird reservoirs will sustain an infectious viremia for 1 to 4 days.

In 2002, evidence of WNV transmission was shown to occur via the transfer of all blood product components including platelets, packed red blood cells, and plasma. Beginning 2003, blood donors were screened for WNV infection utilizing polymerase chain reaction (PCR) testing. Millions of units of blood were screened for WNV utilizing PCR based technology, testing donor mini-pools. Though asymptomatic donors have been identified as positive for WNV in LAC, no transmission associated with blood products has been reported. Additional routes of transmission that have been documented include transplantation of WNV-infected organs, transplacental (mother-to-child), occupational exposures, and through breast milk.



Clinical Infection and Diagnosis

Most persons who become infected with WNV will not develop clinical illness or symptoms. About 20% of persons infected will develop WNV fever with symptoms that include fever, headache, rash, muscle weakness, fatigue, nausea and vomiting, and occasionally lymph node swelling. Approximately one in 150 patients will develop more severe illness, manifesting as WNV neuro-invasive disease (NID). WNV NID includes encephalitis, meningitis, and acute flaccid paralysis (AFP). WNV-associated encephalitis is commonly associated with fever, altered mental status, headache, and seizures; WNV encephalitis usually necessitates high levels of specialized medical care. Focal neurologic deficits, including limb paralysis, cranial nerve palsies, Parkinsonian-like tremors, and other movement disorders have been observed. WNV-associated meningitis usually involves fever, headache, and stiff neck, and has a good prognosis.

DISEASE ABSTRACT




- € The overall incidence of reported WNV infections in 2007 was 0.44 cases per 100,000 population, rising from a low of 0.17 in 2006 when only 16 cases were confirmed (Figure 1).
- € Case fatalities (n=5) occurred for the first time since 2004.
- € Meningitis continues to be the most commonly reported clinical condition, comprising 32% (n=14) of cases.
- € Most WNV infections occurred in persons residing in San Fernando Valley.

STRATIFIED DATA

Trends: WNV infection, including in asymptomatic blood donors, occurred at an incidence rate of 0.44 per 100,000 population in 2007. Both the total number and incidence of WNV infection has decreased dramatically since 2004 when 309 cases were confirmed at an incidence of 3.3 cases per 100,000 population (incidence re-calculated with updated population estimates) (Figure 1).

Seasonality: Onset of cases occurred July through October and peaked in September (Figure 2). Since 2004, the onset of WNV cases has been limited to July through October.

Age: The median age was 62 years (range: 15–94 years). The highest incidence occurred in the 65 and over age group (1.9 per 100,000) (data not shown).





Almost all cases (n=40, 93%) were at least 45 years old.

Sex: Over three times as many male WNV cases were reported than female cases, a rate ratio of 3.4:1. The incidence rates were 0.68 cases and 0.20 cases per 100,000, respectively.

Race/Ethnicity: In 2007, WNV cases occurred only in whites and Hispanics, with whites accounting for the greatest proportion of reported cases (72%) as well as the highest incidence rates of infection (n=31, 1.1 per 100,000). Hispanics comprised 28% of cases (n=12, 0.26 per 100,000) (Figure 3).

Location: The greatest number of reported WNV cases were reported from SPA 2, the San Fernando Valley area (n=27, 1.3 per 100,000). The second highest incidence occurred in SPA 3, the San Gabriel Valley area (n=9, 0.51 per 100,000). WNV occurred sparsely and sporadically in the remaining SPA locations (Figure 4).

Disease Severity: The WNV infections reported presented most frequently as neuroinvasive disease (n=28, 65%); 12 were diagnosed as encephalitis, 14 as meningitis, and 2 as acute flaccid paralysis. A substantial number of infections were asymptomatic blood donors (n=7, 16%). Of those symptomatic cases, 86% (n=31) were hospitalized. Five fatalities (12%) occurred in 2007, the first since 2004 when 14 deaths (5% of cases) were reported. Four of the deaths were diagnosed with encephalitis and one with WNV fever.

COMMENTS

The first symptomatic WNV case in LAC associated with environmental evidence was documented in 2003. In 2004, an outbreak of 309 WNV infections, including asymptomatic blood donors, with 14 deaths were reported in LAC — the most of any CA jurisdiction. In response to the outbreak, LAC DPH added WNV infection to its list of reportable diseases by authority of the Health Officer under California Code of Regulations, Title 17, Sections 2511 and 2505. Physicians and laboratories are required to report all positive laboratory findings of WNV tests to the DPH within one working day.

The following years presented a markedly different picture, with numbers declining to a low of 16 in 2006. This year, however, over twice as many cases were reported. The rise in cases, as well as the continued detection of positive mosquito pools, dead birds and other reservoir animals, has demonstrated that WNV remains endemic in the LAC and southern CA region. As the number of cases has fluctuated greatly from year to year (ranging from 16 to 43 since 2005), the baseline level of cases expected for this region remains to be seen. Sustained surveillance of humans, as well as other animals, will be required in the coming years to help guide public health officials in providing targeted health education to communities at particularly high risk.

PREVENTION

Prevention and control of WNV and other arboviral diseases is most effectively accomplished through integrated vector management programs. These programs include surveillance for WNV activity in mosquito vectors, birds, horses, other animals, and humans; and implementation of appropriate mosquito control measures to reduce mosquito populations when necessary. Additionally, when virus activity is detected in an area, residents are alerted and advised to increase measures to reduce contact with mosquitoes. Currently, there is no human vaccine available against WNV but several vaccines are under development. Important preventive measures against WNV include the following:

- € Apply insect repellent to exposed skin. A higher percentage of DEET in a repellent will provide longer protection. DEET concentrations higher than 50% do not increase the length of protection.
- € When possible, wear long-sleeved shirts and long pants when outdoors for long periods of time.
- € Stay indoors at dawn, dusk, and in the early evening, which are peak mosquito biting times.
- € Help reduce the number of mosquitoes in areas outdoors by draining sources of standing water. This will reduce the number of places mosquitoes can lay their eggs and breed.



A wide variety of insect repellent products are available. CDC recommends the use of products containing active ingredients which have been registered with the U.S. Environmental Protection Agency (EPA) for use as repellents applied to skin and clothing. EPA registration of repellent active ingredients indicates the materials have been reviewed and approved for efficacy and human safety when applied according to the instructions on the label. Of the active ingredients registered with the EPA, three have demonstrated a higher degree of efficacy in the peer-reviewed, scientific literature. Products containing these active ingredients typically provide longer-lasting protection than others:

DEET (N,N-diethyl-m-toluamide)
Picaridin (KBR 3023)
Oil of lemon eucalyptus

Oil of lemon eucalyptus [p.menthane 3, 8-diol (PMD)], a plant based repellent, is registered with EPA. In two recent scientific publications, when oil of lemon eucalyptus was tested against mosquitoes found in the US it provided protection similar to repellents with low concentrations of DEET.

VECTOR CONTROL

There are five local mosquito and vector control districts within LAC that provide mosquito abatement services to all areas of the county. They carry out mosquito and sentinel chicken surveillance, provide public information, and are critical to mosquito-borne disease control. They include:

- € Greater Los Angeles County Vector Control District (GLACVCD)
- € San Gabriel Valley Mosquito and Vector Control District (SGVMVCD)
- € Los Angeles County West Vector Control District (LACWVCD)
- € Antelope Valley Mosquito and Vector Control District (AVMVCD)
- € Compton Creek Mosquito Abatement District (CCMAD)

These five local mosquito and vector control districts work closely with the ACDC to investigate confirmed and presumptive human cases of locally acquired mosquito-borne disease to identify mosquito breeding sites and to put into place appropriate control measures.

ADDITIONAL RESOURCES

- € Centers for Disease Control and Prevention: <http://www.cdc.gov/ncidod/dvbid/westnile/index.htm>
- € California Department of Health Services: <http://www.westnile.ca.gov>
- € Acute Communicable Disease Control Program, Los Angeles County Public Health: <http://www.lapublichealth.org/acd/index.htm>
- € Vector Management Environmental Health, Los Angeles County Public Health: <http://www.lapublichealth.org/eh/index.htm>
- € For additional information on EPA-registered repellents: <http://www.epa.gov/pesticides/factsheets/insectrp.htm>


Mosquito and Vector Control District Websites:

- € Greater Los Angeles County Vector Control District: <http://www.glacvcd.org>
- € West Los Angeles Vector Control District: <http://www.lawestvector.org>
- € San Gabriel Valley Mosquito and Vector Control District: <http://www.sgvmosquito.org>
- € Antelope Valley Mosquito and Vector Control District: <http://www.avmosquito.org>
- € Mosquito and Vector Control Association of California: <http://www.mvcac.org>

WEST NILE VIRUS

CRUDE DATA	
Number of Cases	16
Incidence LAC ^a	
LA County	0.17
California	N/A
United States	N/A
Age at Diagnosis	
Mean	50.9
Median	50.5
Range	28–82 years

^a Cases per 100,000 population.



DESCRIPTION

Life Cycle and Epidemiology

West Nile virus (WNV) is a single-stranded RNA virus placed within the family Flaviviridae, genus Flavivirus. Within the genus Flavivirus, WNV has been serologically classified within the Japanese encephalitis (JE) virus antigenic complex, which includes the human pathogens JE, Murray Valley encephalitis, Saint Louis encephalitis (SLE), and Kunjin viruses.

WNV was indigenous to Africa, Asia, Europe, and Australia, and was introduced to North America in 1999, when it was first detected in New York City. The likely origin of the introduced strain was the Middle East, but the mode of introduction remains unknown. Since 1999, human and non-human WNV surveillance data has documented that WNV has extended its range through most of the continental United States as well as to Canada and Mexico.

The life cycle of the virus involves the transmission of the virus between mosquitoes and bird reservoir hosts. Humans are incidentally infected when bitten by an infected mosquito, usually a *Culex* or *Anopheles* species. The incubation period for human infection is 2 to 14 days. Birds, especially corvids such as the North American crow, are the optimal hosts for harboring and replicating the virus. Mosquitoes become infected when they feed on infected birds, which may circulate high level of viremia for several days. Infectious mosquitoes carry virus particles in their salivary glands and infect susceptible bird species during blood-meal feeding. Bird reservoirs will sustain an infectious viremia for 1 to 4 days. Additional routes of transmission that have been documented include transplantation of WNV-infected organs, blood transfusions, transplacental (mother-to-child), occupational exposures, and through breast milk.


Clinical Infection and Diagnosis

Most persons who become infected with WNV will not develop clinical illness or symptoms. Approximately one in 150 patients will develop more severe illness, manifesting as WNV neuro-invasive disease (NID), and about 20% of persons infected will develop WNV fever with symptoms that include fever, headache, rash, muscle weakness, fatigue, nausea and vomiting, and occasionally lymph node swelling. WNV NID includes encephalitis, meningitis, and acute flaccid paralysis (AFP). WNV-associated encephalitis is commonly associated with the following symptoms: fever, altered mental status, headache, and seizures; WNV encephalitis usually necessitates high levels of specialized medical care. Focal

neurologic deficits, including limb paralysis, cranial nerve palsies, Parkinsonian-like tremors, and other movement disorders have been observed. WNV-associated meningitis usually involves fever, headache, and stiff neck, and has a good prognosis.


DISEASE ABSTRACT

- The overall incidence of reported WNV infections in 2006 was 0.17 cases per 100,000 population, far lower than the incidence rates of previous years, when 3.2 per 100,000 and 0.46 per 100,000 were confirmed in 2004 and 2005, respectively (Figure 1).
- There were no case fatalities in 2005 or 2006.
- Meningitis was the most commonly reported clinical condition as it was in 2005, comprising 25% (n=4) of cases. In 2005, meningitis comprised 34.8% of cases (n=15).
- There were few or no cases in children in both 2005 and 2006.
- Most WNV infections occurred in persons residing in San Fernando Valley.




STRATIFIED DATA

Trends: WNV infection, including in asymptomatic blood donors, occurred at an incidence rate of 0.17 per 100,000 population in 2006. Both the total number and incidence of WNV infection decreased dramatically since 2004 when 309 cases were confirmed at an incidence of 3.2 cases per 100,000 population. In 2005, the incidence was 0.46 per 100,000 (n=43) (Figure 1).



Seasonality: Onset of cases occurred July through October and peaked in August (Figure 2). A similar epidemiologic symptom onset curve also occurred in 2005.

Age: The median age was 50.5 years (range: 28–82 years). For age groups ≥35 years, the incidence rates were similar (they ranged 0.2-0.4 cases per 100,000). There was more varied distribution in 2005 where incidence rates ranged from 0.3 cases per 100,000 among children under 10 to 11.6 cases per 100,000 in those greater than 80 years old.



Sex: A higher proportion of male WNV cases were reported than female cases. The incidence rates were 0.25 cases and 0.08 cases per 100,000, respectively.

Race/Ethnicity: Whites had the greatest proportion of reported cases (81%) as well as the highest incidence rates of infection (n=13, 0.45 per 100,000). Latinos accounted for 13% of cases (n=2, 0.04 per 100,000), and only 6% of reported cases occurred among Asian Pacific Islanders (n=1, 0.1 per 100,000). No cases in

blacks were reported (Figure 3).

Location: The greatest number of reported WNV cases were reported from SPA 2 (n=10, 0.47 per 100,000). WNV cases occurred in only two other areas: SPAs 3 and 4. WNV was distributed more widely in 2005, though SPA 2 also accounted for most cases.

PREVENTION

Prevention and control of WNV and other arboviral diseases is most effectively accomplished through integrated vector management programs. These programs include surveillance for WNV activity in mosquito vectors, birds, horses, other animals, and humans; and implementation of appropriate mosquito control measures to reduce mosquito populations when necessary. Additionally, when virus activity is detected in an area, residents are alerted and advised to increase measures to reduce contact with mosquitoes. Currently, there is no human vaccine available against WNV but several vaccines are under development. Important preventive measures against WNV include the following:

- Apply insect repellent to exposed skin. A higher percentage of DEET in a repellent will provide longer protection. DEET concentrations higher than 50% do not increase the length of protection.
- When possible, wear long-sleeved shirts and long pants when outdoors for long periods of time.
- Stay indoors at dawn, dusk, and in the early evening, which are peak mosquito biting times.
- Help reduce the number of mosquitoes in areas outdoors by draining sources of standing water. This will reduce the number of places mosquitoes can lay their eggs and breed.

A wide variety of insect repellent products are available. CDC recommends the use of products containing active ingredients which have been registered with the U.S. Environmental Protection Agency (EPA) for use as repellents applied to skin and clothing. EPA registration of repellent active ingredients indicates the materials have been reviewed and approved for efficacy and human safety when applied according to the instructions on the label. Of the active ingredients registered with the EPA, three have demonstrated a higher degree of efficacy in the peer-reviewed, scientific literature. Products containing these active ingredients typically provide longer-lasting protection than others:

- DEET (N,N-diethyl-m-toluamide)
- Picaridin (KBR 3023)
- Oil of lemon eucalyptus

Oil of lemon eucalyptus [p.menthane 3, 8-diol (PMD)], a plant based repellent, is registered with EPA. In two recent scientific publications, when oil of lemon eucalyptus was tested against mosquitoes found in the US it provided protection similar to repellents with low concentrations of DEET.

In 2002, evidence of WNV transmission was shown to occur via the transfer of all blood product components including platelets, packed red blood cells, and plasma. Beginning 2003, blood donors were screened for WNV infection utilizing polymerase chain reaction (PCR) testing. Millions of units of blood were screened for WNV utilizing PCR based technology, testing donor mini-pools. Though asymptomatic donors have been identified as positive for WNV in LAC, no transmission associated with blood products has been reported.

COMMENTS

The first symptomatic WNV case in LAC with associated environmental evidence was documented in 2003. In 2004, an outbreak of 309 WNV infections, including asymptomatic blood donors, with 14 deaths were reported in LAC — the most of any CA jurisdiction. The following years have presented a markedly different picture. In 2005, the county only documented 43 infections and no deaths. The decline continued in 2006, during which only 16 cases and no deaths were reported.

In response to the 2004 WNV outbreak, LAC DPH specifically added WNV infection to its list of reportable diseases by authority of the Health Officer under California Code of Regulations, Title 17, Sections 2503 and 2505. Physicians and laboratories are required to report all positive laboratory findings of WNV to the DPH within one working day. Continued vector surveillance efforts have demonstrated that, despite the decline in incidence in LAC, WNV remains endemic (enzootic) in the LAC and southern CA region. Sustained surveillance of humans, as well as other animals, will be required in the coming years to help guide public health officials in providing targeted health education to communities at particularly high risk.

VECTOR CONTROL

There are five local mosquito and vector control districts within LAC that provide mosquito abatement services to all areas of the county. They carry out mosquito and sentinel chicken surveillance, provide public information, and are critical to mosquito-borne disease control. They include:

- Greater Los Angeles County Vector Control District (GLACVCD)
- San Gabriel Valley Mosquito and Vector Control District (SGVVCD)
- Los Angeles County West Vector Control District (LACWVCD)
- Antelope Valley Mosquito and Vector Control District (AVMVCD)
- Compton Creek Mosquito Abatement District

These five local mosquito and vector control districts work closely with the ACDC to investigate confirmed and presumptive human cases of locally acquired mosquito-borne disease to identify mosquito breeding sites and to put into place appropriate control measures.

ADDITIONAL RESOURCES

- Centers for Disease Control and Prevention: www.cdc.gov/ncidod/dvbid/westnile/index.htm
- California Department of Health Services: www.westnile.ca.gov
- Acute Communicable Disease Control Program, Los Angeles County Public Health: www.lapublichealth.org/acd/index.htm
- Vector Management Environmental Health, Los Angeles County Public Health: www.lapublichealth.org/eh/index.htm
- For additional information on EPA-registered repellants: www.epa.gov/pesticides/factsheets/insectrp.htm


Mosquito and Vector Control District Websites:

- Greater Los Angeles County Vector Control District: www.glacvcd.org
- West Los Angeles Vector Control District: www.lawestvector.org
- San Gabriel Valley Mosquito and Vector Control District: www.sgvmosquito.org
- Antelope Valley Mosquito and Vector Control District: www.avmosquito.org
- Mosquito and Vector Control Association of California: www.mvcac.org

WEST NILE VIRUS

CRUDE DATA	
Number of Cases	43
Incidence LAC ^e	
LA County	0.46
California	2.6
United States	1.1
Age at Diagnosis	
Mean	52
Median	56
Range	4-89 years
Case Fatality	0
LA County	0%
California	2%
United States	4%

^a Cases per 100,000 population and based on 2005 population estimates.



DESCRIPTION

LIFE CYCLE AND EPIDEMIOLOGY

West Nile virus (WNV) is a single-stranded RNA virus placed within the family Flaviviridae, genus Flavivirus. Within the genus Flavivirus, WNV has been serologically classified within the Japanese encephalitis (JE) virus antigenic complex, which includes the human pathogens JE, Murray Valley encephalitis, Saint Louis encephalitis (SLE), and Kunjin viruses.

WNV was indigenous to Africa, Asia, Europe, and Australia, and was introduced to North America in 1999, where it was first detected in New York City. The likely origin of the introduced strain was the Middle East, but the mode of introduction remains unknown. From 1999 through 2004, human and non-human WNV surveillance data has documented that WNV has extended its range through most of continental United States to include 47 states and the District of Columbia, Canada, and Mexico. In 2005, 3000 confirmed human WNV cases were reported nationally to the Centers for Disease Control and Prevention (CDC). California (CA) reported the greatest number of any state, 935 cases; Los Angeles County (LAC) reported 43 human cases.

The life cycle of the virus involves the transmission of the virus from a bird reservoir to humans via *Culex*, or *Anopheles* mosquitoes. Birds, especially, corvids such as the North American crow, are the optimal hosts for harboring and replicating the virus. Mosquitoes become infected when they feed on infected birds, which may circulate high level of viremia for several days. Infectious mosquitoes carry virus particles in their salivary glands and infect susceptible bird species during blood-meal feeding. Bird reservoirs will sustain an infectious viremia for 1 to 4 days. Additional routes of transmission that have been documented include transplantation of WNV-infected organs, blood transfusions, transplacental (mother-to-child), occupational exposures, and through breast milk.

CLINICAL INFECTION AND DIAGNOSIS

Most persons who become infected with WNV will not develop clinical illness or symptoms. Approximately one in 150 people will develop more severe illness, manifesting as WNV neuro-invasive disease (NID), and about 20% of persons infected will develop WNV fever with symptoms that include fever, headache, rash, muscle weakness, fatigue, nausea and vomiting, and occasionally lymph node swelling. WNV NID includes encephalitis, meningitis, and acute flaccid paralysis (AFP). WNV-associated encephalitis is commonly associated with the following symptoms: fever, altered mental status, headache, seizures, and usually necessitates high levels of specialized medical care. Focal neurologic deficits, including limb paralysis, cranial nerve palsies, Parkinsonian-like tremors, and movement disorders have also been observed. WNV-associated meningitis usually involves fever, headache, and stiff neck, and has an excellent prognosis.

DISEASE ABSTRACT

- The overall incidence of reported WNV infections including asymptomatic blood donors in 2005 was 0.46 cases per 100,000 population, far lower than the 2004 incidence of 3.2 cases per 100,000 (Figure 1).
- There were no case fatalities in 2005.
- WNV meningitis was the most commonly reported clinical condition, comprising 34.8% (15) of cases in 2005.
- The lowest incidence rates were found in children; there was a consistent increase in incidence after age 40 years.
- Non-Hispanic Whites had the highest proportion of symptomatic infections (49%, n=21) followed by Hispanics (40%, n=17) (Figure 2).
- Most WNV infections occurred in persons residing in San Fernando Valley.

STRATIFIED DATA


Trends: In 2003, the first WNV case with associated environmental evidence was documented in LAC. In 2004, 209 cases were document in LAC, the most of any CA jurisdiction. In 2005, both the total number of WNV cases and the incidence of WNV infection decreased dramatically with 43 cases reported, an incidence of 0.46 cases per 100,000 population.

Seasonality: Cases were reported from the beginning of summer (first week of July) through fall (last week of October). The peak onset of cases was the 2nd week in August (33 week) when 9 cases were reported (Figure 1).

Age: The median age for all WNV cases including asymptomatic blood donors was 43 years old (range: 18-57 years). The median age for both WNF and WNV meningitis was 47 years. The median age for the 13 encephalitis cases was 63 years old. The lowest incidence rates of WNV infection occurred among children under 10, 0.3 cases per 100,000, whereas the greatest incidence occurred in those > 80 years (11.6 cases per 100,000).

Sex: A higher proportion of male WNV cases were reported than female cases. The incidence rates were 0.5 cases and 0.4 cases per 100,000, respectively.

Race/Ethnicity: Whites had the both the greatest proportion of reported cases and highest incidence rates of infection, 49% of cases (n=21, 0.7 per 100,000), followed by Latinos, 40% of cases (n= 17, 0.4 per 100,000). Only 5% of reported cases occurred among Asian Pacific Islanders (n=2, 0.2 per 100,000) and only 1 case (1%) was reported in an African-American (0.1 per 100,000) (Figure 2).



Location: The greatest number of reported WNV cases were reported from SPA 2 (n=19, 0.9 per 100,000). Whereas, SPA 7 (n=12, 4.0 per 100,000) had the second largest number of reported cases but had the highest incidence WNV infection (Figure 3) of any of the service planning areas.

PREVENTION

Prevention and control of WNV and other arboviral diseases is most effectively accomplished through integrated vector management programs. These programs include surveillance for WNV activity in mosquito vectors, birds, horses, other animals, and humans, and implementation of appropriate mosquito control measures to reduce mosquito populations when necessary. Additionally, when virus activity is detected in an area, residents are alerted and advised to increase measures to reduce contact with mosquitoes. Currently, there is no human vaccine available against WNV but several vaccines are under development. Important preventive measures against WNV include the following:

- Apply insect repellent to exposed skin. A higher percentage of DEET in a repellent will provide longer protection. DEET concentrations higher than 50% do not increase the length of protection.
- When possible, wear long-sleeved shirts and long pants when outdoors for long periods of time.
- Stay indoors at dawn, dusk, and in the early evening, which are peak mosquito biting times.
- Help reduce the number of mosquitoes in areas outdoors by draining sources of standing water. This will reduce the number of places mosquitoes can lay their eggs and breed.

A wide variety of insect repellent products are available. CDC recommends the use of products containing active ingredients which have been registered with the U.S. Environmental Protection Agency (EPA) for use as repellents applied to skin and clothing. EPA registration of repellent active ingredients indicates the materials have been reviewed and approved for efficacy and human safety when applied according to the instructions on the label. Of the active ingredients registered with the EPA, three have demonstrated a higher degree of efficacy in the peer-reviewed, scientific literature. Products containing these active ingredients typically provide longer-lasting protection than others:


- DEET (N,N-diethyl-m-toluamide)
- Picaridin (KBR 3023)
- Oil of lemon eucalyptus

Oil of lemon eucalyptus [p.menthane 3, 8-diol (PMD)], a plant based repellent, is also registered with EPA. In two recent scientific publications, when oil of lemon eucalyptus was tested against mosquitoes found in the US it provided protection similar to repellants with low concentrations of DEET.

In 2002, WNV transmission was documented from all components of blood products including platelets, packed red blood cells, and plasma. Subsequently in 2003, all blood donors were screened for WNV infection utilizing polymerase chain reaction (PCR) testing. Millions of units of blood were screened for WNV utilizing PCR based technology, testing donor mini-pools. In 2005, WNV screening continued and no transmission associated with blood products were reported.

COMMENTS

In 2005, 43 human WNV cases were confirmed in LAC among hundreds of tested patients. WNV is considered endemic (enzootic) to LAC and Southern California. Sustained surveillance will be required in



the coming years, including surveillance among humans, dead birds, mosquito pools, and sentinel chickens. These activities guide public health officials in providing targeted health education to communities at particularly high risk. In response to the 2004 WNV outbreak, LAC DHS specifically added WNV infection to its list of reportable diseases by authority of the Health Officer under California Code of Regulations, Title 17, Sections 2503 and 2505. Physicians and laboratories are required to report all positive laboratory findings of WNV to the Department of Health Services within one (1) working day.

VECTOR CONTROL

There are five local mosquito and vector control districts within LAC that provide mosquito abatement services to all areas of the county. They carry out mosquito and sentinel chicken surveillance, provide public information, and are critical to mosquito-borne disease control. They include:

- Greater Los Angeles County Vector Control District (GLACVCD)
- San Gabriel Valley Mosquito and Vector Control District (SGVVCD)
- Los Angeles County West Vector Control District (LACWVCD)
- Antelope Valley Mosquito and Vector Control District (AVMVCD)
- Compton Creek Mosquito Abatement District

These five local mosquito and vector control districts work closely with the ACDC to investigate confirmed and presumptive human cases of locally acquired mosquito-borne disease to identify mosquito breeding sites and to put into place appropriate control measures.

REFERENCES:

1. Campbell GL, Marfin AA, Lanciotti RS, Gubler DJ. West Nile virus. *Lancet Infect Dis* 2002 ;519-29.
2. Watson JT, Pertel PE, Jones RC, et al. Clinical characteristics and functional outcomes of West Nile fever. *Ann Intern Med.* 2004;141:360-365.
3. Sejvay JJ, Bode AV, Marfin AA, Campbell, et al. West Nile virus-associated flaccid paralysis. *EID* 2005; 11:7:1021-1026.
4. Somomon T. Flavivirus Encephalitis. *N Engl J Med* 2004;351:370-8.
5. Reisen W, Lothrop H, Chiles R, et al. West Nile virus in California. *EID* 2004; 10:8: 1369-1377.
6. Nash D, Mostasharia F, Fine A, Miller M, et al. The outbreak of West Nile virus infection in the New York City area in 1999. *N Eng J Med* 2001; 344:24: 1807-1814.

ADDITIONAL RESOURCES

- Centers for Disease Control and Prevention: www.cdc.gov/ncidod/dvbid/westnile/index.htm
- California Department of Health Services: www.westnile.ca.gov/.
- Acute Communicable Disease Control Program, Los Angeles County Public Health: www.lapublichealth.org/acd/index.htm/
- Vector Management Environmental Health, Los Angeles County Public Health: www.lapublichealth.org/eh/index.htm/
- For additional information on EPA-registered repellants: www.epa.gov/pesticides/factsheets/insectrp.htm

Mosquito and Vector Control District Websites:


- Greater Los Angeles County Vector Control District: www.glacvcd.org/
- West Los Angeles Vector Control District: www.lawestvector.org/
- San Gabriel Valley Mosquito and Vector Control District: www.sgvmosquito.org/
- Antelope Valley Mosquito and Vector Control District: www.avmosquito.org/
- Mosquito and Vector Control Association of California: www.mvcac.org/



WEST NILE VIRUS

CRUDE DATA	
Number of Cases	309
Incidence LAC ^a	
LA County	3.2
California	0.81
United States	0.39
Age at Diagnosis	
Mean	54
Median	54
Range	<5–94 years
Case Fatality	14
LA County	4.5%
California	3.6%
United States	3.9%

^a Cases per 100,000 population and based on 2004 population estimates.



DESCRIPTION

LIFE CYCLE AND EPIDEMIOLOGY

West Nile virus (WNV) is a single-stranded RNA virus placed within the family Flaviviridae, genus Flavivirus. Within the genus Flavivirus, WNV has been serologically classified within the Japanese encephalitis (JE) virus antigenic complex, which includes the human pathogens JE, Murray Valley encephalitis, Saint Louis encephalitis (SLE), and Kunjin viruses.

WNV is indigenous to Africa, Asia, Europe, Middle East, and Australia, and was introduced to North America in 1999, where it was first detected in New York City. The likely origin of the introduced strain was the Middle East, but the mode of introduction remains unknown. From 1999 through 2004, human and non-human WNV surveillance data has documented that WNV has extended its range through most of continental United States to include 47 states and the District of Columbia, Canada, and Mexico. In 2004, 2539 confirmed human WNV cases were reported nationally to the Centers for Disease Control and Prevention (CDC). California (CA) reported the greatest number of any state, 779 cases; Los Angeles County (LAC) reported 309 human cases.

The life cycle of the virus involves the transmission of the virus from a bird reservoir to humans via *Culex*, or Anopheles mosquitoes. Birds, especially, corvids such as the *North American crow*, are the optimal hosts for harboring and replicating the virus. Mosquitoes become infected when they feed on infected birds, which may circulate high level of viremia for several days. Infectious mosquitoes carry virus particles in their salivary glands and infect susceptible bird species during blood-meal feeding. Bird reservoirs will sustain an infectious viremia for 1 to 4 days. Additional less frequent routes of transmission that have been documented include transplantation of WNV-infected organs, blood transfusions, transplacental (mother-to-child), occupational exposures, and through breast milk.



CLINICAL INFECTION AND DIAGNOSIS

Most persons who become infected with West Nile virus (WNV) will not develop clinical illness or symptoms. The incubation period for WNV infection can range from 2 to 14 days, although longer incubation periods have been documented in immunosuppressed persons. Approximately one in 150 people will develop severe illness, WNV neuro-invasive disease (NID), and about 20% of persons infected with WNV will develop WNV fever (WNVF) with symptoms that can include fever, headache, rash, muscle weakness, fatigue, nausea and vomiting, and occasionally lymph node swelling. WNV NID includes: encephalitis, meningitis, and acute flaccid paralysis (AFP). WNV-associated encephalitis, the most severe form of NID, is commonly associated with the following symptoms: fever, altered mental status, headache, seizures, and usually necessitates high levels of specialized medical care. Focal neurologic deficits, including limb paralysis, cranial nerve palsies, Parkinsonian-like tremors, and movement disorders have also been observed. WNV-associated meningitis usually involves fever, headache, and stiff neck. WNV-associated poliomyelitis or acute flaccid paralysis (AFP) syndrome, is the least common among NID. AFP, initially described as atypical Guillian Barre Syndrome in New York City in 1999, was well documented in 2002 during the outbreak in Louisiana. Clinically this syndrome is characterized by the acute onset of asymmetric limb weakness or paralysis in the absence of sensory loss. The paralysis can occur in the absence of fever, headache, or other common symptoms associated with WNV infection. AFP is associated with significant short- and long-term illness and death.

WNV infection is suspected in a person based on clinical symptoms and patient history. Serologic laboratory testing is required to confirm a diagnosis. Diagnosis of acute infection requires the detection of IgM antibody. Serum IgM is usually positive within 5 to 14 days of illness in over 90% of cases and CSF IgM is positive within 7 days of onset. The most commonly performed serologic test is the IgM antibody-capture assay, the enzyme-linked immunosorbent assay (MAC-ELISA). Los Angeles County Public Health laboratory utilizes both IgM and IgG antibody-immunofluorescent Assay (IFA) as well as IgM enzyme immunoassays (EIA). The plaque-reduction neutralization test (PRNT), a cell culture based assay, is a confirmatory test performed at the CA Department of Health Services (DHS) Viral and Rickettsial Disease Laboratory (VRDL). PRNT can distinguish between arthropod-borne flaviviruses such as SLE and WNV. It can also be used to help distinguish false-positive results in an IgM antibody-capture enzyme-linked immunosorbent assay or other assays. In addition to PRNT and serologic testing, nucleotide based testing such as PCR is utilized for blood screening and for surveillance of WNV-infected mosquitoes and dead birds.

HISTORY OF WNV IN LOS ANGELES COUNTY

In 2002, a single human case of WNV-associated meningitis was confirmed. The young woman recovered uneventfully and denied history of travel outside of LAC, blood transfusions, or a history of organ transplantation. However, in 2002, there was no environmental evidence documenting the presence of WNV within LAC such as WNV-infected dead birds, mosquito pools or sentinel chickens with WNV sero-conversion. The first environmental evidence that WNV had arrived to LAC occurred in the summer of 2003 with the presence of WNV-infected dead birds, mosquito pools, and sero-positive sentinel chickens. In 2003, one human case of WNVF, with symptom onset in mid-October, was laboratory confirmed, and was most likely infected by local WNV-infected mosquitoes. In June 2004, the LAC public health department documented the first of many human cases.

DISEASE ABSTRACT

- The overall incidence of reported WNV infections including asymptomatic blood donors in 2004 was 3.2 cases per 100,000 population (Figure 1).
- WNVF was the most common WNV-associated clinical condition, 149 cases or 48.2 % of reported cases in 2004.
- There were 14 case fatalities, including 10 with encephalitis, 2 with meningitis, and 2 WNVF cases. The mean age was 76.4 years (range 60-94).
- NID and deaths were each associated with older age.
- The lowest incidence of WNV infection was found in children; there was a dramatic increase in WNV




incidence after age 40 years (Figure 2).

- Non-Hispanic Whites had the highest proportion of symptomatic infections (57%) followed by Hispanics (39%).
- Significantly more male cases (65%, n=200) were reported than female (35%, n=109) in 2004.
- Most WNV infections occurred in persons residing in suburban valleys, areas close to the San Gabriel River and hillside communities.

STRATIFIED DATA

Seasonality: Cases were reported from late spring (first week of June) through fall (second week of October). The peak onset of cases was the 2nd week in August (week 33) when 41 cases were recorded (Figure 1).

Age: The median age for all reported WNV cases including asymptomatic blood donors was 54 years old (range: 5–94 years). Of the 149 WNF cases, the median age was 51 years (range: 6–91). The median age for the 82 meningitis cases was 53 years (range: 5–90). The median age for the 47 encephalitis cases was 70 years old (range: 28–94). The median age for the 7 reported AFP was 45 years (range: 34–78). The median age for the 14 fatal cases was 77.0 years (range: 60–94). The incidence of WNV infection increased steadily with age in Whites and Hispanics (Figure 2). The lowest incidence rates occurred in children under 10, 0.3 cases per 100,000, whereas the greatest incidence occurred in those > 80 years (11.6 cases per 100,000).




Sex: Males were almost twice as likely to present with WNV infection compared to females. The overall LAC WNV incidence rate was 4.2 cases per 100,000 male population versus 2.3 cases per 100,000 female population.

Race/Ethnicity: Whites had the both the greatest proportion of reported cases and overall highest incidence rates of infection, 57% of cases (n=170, 5.9 per 100,000), followed by Hispanics, 39% of cases (n= 117, 2.6 per 100,000). When incidence rates were reviewed by age group and race/ethnicity, whites had the highest incidence of infection in individuals age 30 and above, whereas, Hispanics had the highest incidence in age groups under age 20 years (Figure 2). Only 3% of reported cases occurred among Asian Pacific Islanders (n=8, 0.6 per 100,000) and 1% among Blacks (n=3, 0.3 per 100,000).

Location: The number of reported WNV cases was highest in SPA 3 (n=114, 6.7 per 100,000), SPA 2 (n=84, 4.0 per 100,000), and SPA 7 (n=77, 5.6 per 100,000) (Figure 3).

PREVENTION

Prevention and control of WNV and other arboviral diseases is most effectively accomplished through integrated vector management programs. These programs include surveillance for WNV activity in mosquito vectors, birds, horses, other animals, and humans, and implementation of appropriate mosquito control measures to reduce mosquito populations when necessary. Additionally, when virus activity is detected in an area, residents are alerted and advised to increase measures to reduce contact with





mosquitoes. Currently, there is no human vaccine available against WNV but several vaccines are under development. Important preventive measures against WNV include the following:

- Applying insect repellent to exposed skin. A higher percentage of DEET in a repellent will provide longer protection. DEET concentrations higher than 50% do not increase the length of protection.
- When possible, wear long-sleeved shirts and long pants when outdoors for long periods of time.
- Staying indoors at dawn, dusk, and in the early evening, which are peak mosquito biting times.
- Help reduce the number of mosquitoes in areas outdoors by draining sources of standing water. This will reduce the number of places mosquitoes can lay their eggs and breed.

A wide variety of insect repellent products are available. CDC recommends the use of products containing active ingredients which have been registered with the U.S. Environmental Protection Agency (EPA) for use as repellents applied to skin and clothing. EPA registration of repellent active ingredients indicates the materials have been reviewed and approved for efficacy and human safety when applied according to the instructions on the label. Of the active ingredients registered with the EPA, two have demonstrated a higher degree of efficacy in the peer-reviewed, scientific literature. Products containing these active ingredients typically provide longer-lasting protection than others:

- DEET (N,N-diethyl-m-toluamide)
- Picaridin (KBR 3023)

Picaridin is an EPA approved mosquito repellent that will be commercially available in 2005.

In 2002, WNV transmission was documented from all components of blood products including platelets, packed red blood cells, and plasma. Subsequently in 2003, all blood donors were screened for WNV infection utilizing PCR testing. Millions of units of blood were screened for WNV utilizing PCR based technology, testing donor mini-pools. Over 1000 donor units were held from distribution in 2003, and only 6 clinical WNV cases were documented to be associated with WNV infected transfusion. In 2004, WNV screening intensified and individual donors were screened in order to detect an even lower level of donor viremia than what could be detected from mini-pool screening.

COMMENTS

In 2004, 309 human WNV cases were confirmed in LAC among hundreds of tested patients. WNV is now considered endemic (enzootic) to LAC and Southern California. Sustained surveillance will be required in the coming years, including surveillance among humans, dead birds, mosquito pools, and sentinel chickens. These activities guide public health officials in providing targeted health education to communities at particularly high risk. In response to the 2004 WNV outbreak, LAC DHS specifically added WNV infection to its list of reportable diseases by authority of the Health Officer under California Code of Regulations, Title 17, Sections 2503 and 2505. Physicians and laboratories are required to report all positive laboratory findings of WNV to the Department of Health Services within one (1) working day.

Medical providers play a key role in providing WNV health education to their patients at high risk. A county-wide phone survey in September 2004 revealed that community knowledge of WNV as a potential health risk in LAC was high, 93%, and most people knew that WNV was transmitted by the bite of a mosquito, 97%. But the survey also showed the public's self-reported change in their behaviors to protect themselves against WNV was at about 50% of persons surveyed. Only 20% of respondents reported using repellent prior to 2004. Promotion of preventive measures can help to minimize the risk of being exposed to WNV is a critical part of the public health message.

VECTOR CONTROL

There are five local mosquito and vector control districts within LAC that provide mosquito abatement services to all areas of the county. They carry out mosquito and sentinel chicken surveillance, provide public information, and are critical to mosquito-borne disease control. They include:

- Greater Los Angeles County Vector Control District (GLACVCD)
- San Gabriel Valley Mosquito and Vector Control District (SGVVCD)
- Los Angeles County West Vector Control District (LACWVCD)



- Antelope Valley Mosquito and Vector Control District (AVMVCD)
- Compton Creek Mosquito Abatement District

These five local mosquito and vector control districts work closely with the ACDC to investigate confirmed and presumptive human cases of locally acquired vector-borne disease to determine the source and conditions of transmission.

GLACVCD is the largest vector control district in LAC serving 4.5 million residents in a 1,330 square mile area covering cities from San Fernando Valley, Los Angeles River, the proximate cities of Maywood, Bell, Huntington Park and portions of LAC. The West Vector Control District covers approximately 600 square miles, contains 23 cities and unincorporated territories of the County of Los Angeles, and provides services for 2,866,000 people. The District includes the cities of Agoura Hills, Beverly Hills, Calabasas, Culver City, El Segundo, Hawthorne, Hermosa Beach, Hidden Hills, Inglewood, Lawndale, Lomita, the westerly portion of LAC, Malibu, Manhattan Beach, Palos Verdes Estates, Rancho Palos Verdes, Redondo Beach, Rolling Hills, Rolling Hills Estates, Santa Monica, Torrance, West Hollywood, Westlake Village, and unincorporated territory of the County of Los Angeles. The San Gabriel Valley Mosquito and Vector Control District covers many parts of the San Gabriel Valley, including cities of Alhambra to West Covina. This district had the most cases of WNV in 2004. The Antelope Valley Mosquito and Vector Control District offers programs that provide information and education for the Antelope Valley general public, schools and community organizations; propagated through brochures, pamphlets, seminars, speeches and presentations. The District encompasses an area of approximately 230 square miles and serves residents within District boundaries in Palmdale, Lancaster and Quartz Hill.

Since mosquitoes serve as vectors for disease transmission, WNV-positive mosquito pools are another critical environmental indicator; as such, mosquito pools are routinely tested for the presence of WNV. In 2004, 378 mosquito pools tested positive in LAC, nearly one-third of the identified positive mosquito pools in California (1,136 pools). The last positive mosquito pool in LAC was identified on October 21, 2004 from Harbor City.

In an effort to help protect the public health from the threat of WNV disease, the LAC DHS funded a one year agreement with local mosquito and vector control districts to provide mosquito abatement services to all areas of the county not currently within the jurisdiction of a control district. At least 186,000 persons reside in such regions. On July 2004, city managers in La Cañada-Flintridge, South Pasadena, and Baldwin Park—cities without mosquito control programs—were notified that the County would provide temporary mosquito abatement services by contracting with the San Gabriel Valley Mosquito and Vector Control District for fiscal year 2004-2005. Because portions of the cities of Palmdale and Lancaster as well as unincorporated Antelope Valley and Santa Clarita Valley regions are also without abatement services, we entered into a similar one-year agreement with the Antelope Valley Mosquito and Vector Control District to provided abatement and surveillance service, pending a permanent solution. The challenge of the coming months will be to follow-up with these cities to ensure that they have plans to continue funding of local mosquito abatement services when the county contracts expire in July 2005.

REFERENCES:

1. Campbell GL, Marfin AA, Lanciotti RS, Gubler DJ. West Nile virus. *Lancet Infect Dis* 2002 ;519-29.
2. Watson JT, Pertel PE, Jones RC, et al. Clinical characteristics and functional outcomes of West Nile fever. *Ann Intern Med.* 2004;141:360-365.
3. Sejvay JJ, Bode AV, Marfin AA, Campbell, et al. West Nile virus-associated flaccid paralysis. *EID* 2005; 11:7:1021-1026.
4. Somomon T. Flavivirus Encephalitis. *N Engl J Med* 2004;351:370-8.
5. Reisen W, Lothrop H, Chiles R, et al. West Nile virus in California. *EID* 2004; 10:8: 1369-1377.
6. Nash D, Mostasharia F, Fine A, Miller M, et al. The outbreak of West Nile virus infection in the New York City area in 1999. *N Eng J Med* 2001; 344:24: 1807-1814.



ADDITIONAL RESOURCES

- Centers for Disease Control and Prevention: www.cdc.gov/ncidod/dvbid/westnile/index.htm
- California Department of Health Services: www.westnile.ca.gov/.
- Acute Communicable Disease Control Program, Los Angeles County Public Health: www.lapublichealth.org/acd/index.htm/
- Vector Management Environmental Health, Los Angeles County Public Health: www.lapublichealth.org/eh/index.htm/
- For additional information on EPA-registered repellants: www.epa.gov/pesticides/factsheets/insectrp.htm

Mosquito and Vector Control District Websites:

- Greater Los Angeles County Vector Control District: www.glacvcd.org/
- West Los Angeles Vector Control District: www.lawestvector.org/
- San Gabriel Valley Mosquito and Vector Control District: www.sgvmosquito.org/
- Antelope Valley Mosquito and Vector Control District: www.avmosquito.org/
- Mosquito and Vector Control Association of California: www.mvacac.org/



**WEST NILE VIRUS WITHIN LOS ANGELES COUNTY:
FIRST AUTOCHTHONOUS HUMAN WEST NILE VIRUS INFECTION IN 2003**

Since the introduction of West Nile virus (WNV) in the continental US in summer of 1999, WNV has become established in nearly all of the contiguous states causing nearly 10,000 cases and 262 deaths in 2003.

In 2002, the first locally acquired human case of WNV in California was identified in Los Angeles County (LAC). However, other forms of local WNV surveillance including dead birds, sentinel chickens, and mosquito pools did not reveal evidence of WNV. In 2003, one case of WNV fever acquired within LAC was documented and laboratory-confirmed.

The first case of locally acquired WNV fever in a LAC resident was laboratory-confirmed in late December 2003 by the California Department of Health Services Viral and Rickettsial Diseases Laboratory (VRDL). ACDC was notified by ARUP Laboratory of Utah of a positive WNV serum IgM test in late October 2003 in a LAC resident. The patient and his physician were contacted and interviewed by ACDC, and the patient provided another serum specimen in early December 2003 so that confirmatory testing could be completed at the LAC Public Health Laboratory.

The case, a 61 year-old Hispanic male, was admitted to an LAC hospital for complaints of fever, fatigue, nausea and diarrhea for 10 days in mid-October 2003. Serum WNV testing was ordered by the attending physician as part of a fever work-up during his hospital admission. The patient recovered uneventfully. He lived in Whittier and gave a history of mosquito bites two days prior to admission while sleeping in his living room with a broken screen door. He believes he was bitten in the early morning of the first week of October. He denied any travel outside of Whittier area 14 days before onset of symptoms. He received no blood products or organ donations within the month prior to symptom onset. In early December, the patient's serum tested weakly positive by CDC WNV ELISA testing in both the LAC Public Health Laboratory and the state VRDL. WNV confirmatory testing, plaque reduction neutralization test, performed at the VRDL confirmed the diagnosis of WNV fever in late December 2003.

Related environmental findings in the fall of 2003 include: 11 dead crows with WNV recovered in Whittier and 6 WNV-infected mosquito pools in two adjoining cities in late September to November 2003. In 2003, there were no sentinel chickens with WNV-positive blood tests from LAC. This is the third endemic case of WNV infection acquired in the state of California in 2003 with the first two human cases being reported from Imperial and San Bernardino counties respectively.